A New Version of the Second Law of Information Thermodynamics Using Entanglement Measure

Hiroyasu Tajima[1]

[1] *Department of Physics, The University of Tokyo, Komaba, Meguro, Tokyo 153-8505*

We present a new version of the second law of information thermodynamics using an entanglement measure as the information content. In the new version $W_{\rm ext}\le -\Delta F+k_BTI_{\rm E}$, the new information $I_{\rm E}$ is the amount of the entanglement taken from the system through a unitary interaction between the system and the probe. The upper bound of the new second law is always achievable in the approximation such that the upper bound of the standard second law $W_{\rm ext}\le -\Delta F$ is always achievable. We also prove that when the system is a two-level system and when we perform only thermodynamic operations such that the equality of $W_{\rm ext}\le -\Delta F$ is achieved, we can always achieve the new upper bound without the above approximation, whereas under the same assumption as above there exists a measurement for which the upper bound of $W_{\rm ext}\le -\Delta F+k_BTI_{\rm QC}$ [6] can*not* be achieved irrespective of the feedback.

arXiv:1212.0407v1 [quant-ph] 3 Dec 2012

The second law of thermodynamics appears to be violated in thermodynamic processes that include measurements and feedbacks; this well-known fact has been the center of attention and numerous studies have long been conducted on such processes [1]–[12]. The second law of information thermodynamics [6] is a monumental landmark of these studies; in the case of an isothermal process, it is expressed as

$$W_{\rm ext}\le -\Delta F+k_BTI_{\rm QC}, \tag{1}$$

where $I_{\rm QC}$ is the QC-mutual information content [6]. This inequality gives a new upper bound for the work extracted from a thermodynamic system when measurement and feedback are permitted on the system. The upper bound is always achievable in the approximation such that the upper bound of the standard second law $W_{\rm ext}\le -\Delta F$ is always achievable [7].

In a different context, the entanglement [17]–[18] has been studied as a resource for various types of information processing [13]–[16]. In recent years, the entanglement has appeared in various fields of physics [19]–[21]. However, the relation between the entanglement and thermodynamic processes with measurements and feedback controls is thus far not known.

In the present Letter, we derive a new version of the second law of information thermodynamics, for which the "information" $I_{\rm E}$ can be expressed in the form of an entanglement measure. In the case of an isothermal process, the second law can be expressed as

$$W_{\rm ext}\le -\Delta F+k_BTI_{\rm E}, \tag{2}$$

where $I_{\rm E}$, which we will define later, represents "the amount of the entanglement between the system and the response system, which the probe takes from the system." This inequality has the following three features. First, we can always achieve the new upper bound in the approximation such that the upper bound of the standard second law $W_{\rm ext}\le -\Delta F$ is always achievable. Second, when the system is a two-level system and when we perform only thermodynamic operations such that the equality of $W_{\rm ext}\le -\Delta F$ is achieved, we can always achieve the new upper bound without the above approximation, through proper interpretation of information and proper feedback. Note that the second feature is not valid for (1); as shown later in this Letter, under the same assumption as above, there exists a measurement for which the upper bound of (1) can*not* be achieved irrespective of the feedback. Third, the quantity $I_{\rm E}$ depends only on the premeasurement state of the system and the unitary interaction between the probe and the system. This implies that the information gain is already completeded when the unitary interaction is over and the projective measurement is only the interpretation of the information.

As the setup, we consider a thermodynamic system S that is in contact with heat baths B_m for $m=1,2,...,n$ which are at temperatures T_1,c,T_n, respectively. We refer to the whole set of baths $\{B_m\}$ as B. Except when we perform measurement or feedback control, we express the Hamiltonian of the whole system as

$$\hat{H}(\vec{\lambda}(t))=\hat{H}^S(\vec{\lambda}^S(t))+\sum_{m=1}^{n}[\hat{H}^{SB_m}(\vec{\lambda}^{SB_m}(t))+\hat{H}^{B_m}], \tag{3}$$

where $\hat{H}^S(\vec{\lambda}^S(t))$ is the Hamiltonian of the system S, $\hat{H}^{B_m}$ is the Hamiltonian of the bath B_m, and $\hat{H}^{SB_m}(\vec{\lambda}^{SB_m}(t))$ is the interaction Hamiltonian between the system S and the heat bath B_m. The Hamiltonian is controlled through external parameters $\vec{\lambda}^S(t)$ and $\vec{\lambda}^{SB}(t)$. We assume that there exists a value of $\vec{\lambda}^{SB}(t)=\vec{\lambda}_0$ such that $\hat{H}^{SB_m}(\vec{\lambda}_0)=\hat{0}$. We call the time evolution of the whole system with controlled $\vec{\lambda}^S(t)$ and $\vec{\lambda}^{SB}(t)$ a thermodynamic operation. Further we assume that we can realize a thermodynamic equilibrium state at temperature T_m by connecting S and B_m and waiting. Note that the equilibrium state may not be a canonical distribution. We define the energy U of a state ρ as ${\rm tr}[\rho\hat{H}]$ and define the Helmholtz free energy F for an equilibrium state at a temperature T as $-k_BT\log Z(\beta)$, where $\beta\equiv(k_BT)^{-1}$ and $Z(\beta)\equiv{\rm tr}[\exp(-\beta\hat{H})]$. A thermodynamic operation from a canonical distribution to an equilibrium state achieves the equality of the following inequality if and only if the final state is a canonical distribution too [6];

$$\frac{U-F}{T}+\sum_{m=1}^{n}\frac{Q_m}{T_m}\le\frac{U'-F'}{T'}, \tag{4}$$

where Q_m is the heat moving from B_m to S during the op-


$t=t_{\rm i}$
thermodynamics process $U_{\rm init}$
$t=t_1$
interaction U_{SP}
$t=t'_1$
$I_{\rm E}$ is determined here
projective measurement $\{P_{(k)}\}$ on probe
$t=t_2$
$I_{\rm QC}$ is determined here
feedback control $\{U_{(k)}\}$ on system and baths
$t=t_3$
thermodynamics Process $U_{\rm fin}$
$t=t_{\rm f}$

FIG. 1: Schematic of the thermodynamic processes from $t_{\rm i}$ to $t_{\rm f}$.

eration, U, F, T are the energy, the free energy and the temperature of the initial state, respectively, and U', F', T' are those of the final state, respectively. We refer to the thermodynamic operation whose initial and final states are the equilibrium states and which achieves the equality of (4) as a strictly quasi-static operation.

Under the above setup, we consider the following thermodynamic process from $t = t_{\rm i}$ to $t = t_{\rm f}$ (Fig. 1). At $t = t_{\rm i}$, we start the process with the following initial state:

$$\rho_{\rm i} = \frac{\exp[-\beta \hat{H}^S_{\rm i}]}{Z^S_{\rm i}(\beta)} \otimes \frac{\exp[-\beta_1 \hat{H}^{B_1}]}{Z^{B_1}(\beta_1)} \otimes ... \otimes \frac{\exp[-\beta_n \hat{H}^{B_n}]}{Z^{B_n}(\beta_n)}, \quad (5)$$

where $\hat{H}^S_{\rm i} = \hat{H}^S(\vec{\lambda}^S(t_{\rm i}))$, $Z^S_{\rm i} = {\rm tr}[\exp[-\beta \hat{H}^S_{\rm i}]]$, $\beta_m = (k_B T_m)^{-1}$ and $Z^{B_m} = {\rm tr}[\exp[-\beta_m \hat{H}^{B_m}]]$ $(m = 1, ..., n)$. From $t = t_{\rm i}$ to $t = t_1$, we perform a thermodynamic operation, which is expressed as $\hat{U}_{\rm init}$. At $t = t_1$, the state is given by $\rho_1 = \hat{U}_{\rm init} \rho_{\rm i} \hat{U}^\dagger_{\rm init}$. Adding a proper reference system R, we can find a pure state $|\psi_{SBR}\rangle$, which satisfies

$${\rm tr}_R[|\psi_{SBR}\rangle \langle \psi_{SBR}|] = \rho_1. \quad (6)$$

From $t = t_1$ to $t = t'_1$, we introduce a unitary interaction U_{SP} between the system S and the probe P, which is initialized to a state $|0_P\rangle$. At $t = t'_1$, the state of the whole system is expressed as

$$|\psi_{PSBR}\rangle = (U_{SP} \otimes \hat{1}_B \otimes \hat{1}_R)(|0_P\rangle \otimes |\psi_{SBR}\rangle), \quad (7)$$

where $\hat{1}_B$ and $\hat{1}_R$ are the identity operators.

At this point, we define a new quantity $I_{\rm E}$, which we call the entanglement information, as follows:

$$\begin{aligned} I_{\rm E} &\equiv E_F^{SB\text{-}R}(|\psi_{SBR}\rangle \langle \psi_{SBR}|) - E_F^{SB\text{-}R}(\rho_{SBR}) \quad (8) \\ &= S(\rho_1) - E_F^{SB\text{-}R}(\rho_{SBR}), \end{aligned}$$

where $S(\rho) \equiv -{\rm tr}[\rho \log \rho]$, $\rho_{SBR} \equiv {\rm tr}_P[|\psi_{PSBR}\rangle \langle \psi_{PSBR}|]$, and $E_F^{SB\text{-}R}(\rho)$ is the entanglement of formation [18] between SB and R;

$$E_F^{SB\text{-}R}(\rho) \equiv \min_{\rho_{SBR}=\sum q_j |\phi^j\rangle\langle\phi^j|} \sum_j q_j E^{SB\text{-}R}(|\phi^j\rangle) \quad (9)$$

with $E^{SB\text{-}R}(|\phi^j\rangle)$ being the entanglement entropy [17] between SB and R for a pure state $|\phi^j\rangle$. Note that $E_F^{SB\text{-}R}(|\psi_{SBR}\rangle \langle \psi_{SBR}|)$ and $E_F^{SB\text{-}R}(\rho_{SBR})$ indicate the amount of entanglement between SB and R at $t = t_1$ and $t = t'_1$, respectively. Thus, $I_{\rm E}$ is the amount of entanglement between SB and R that is taken by the probe P during the interaction U_{SP}.

From $t = t'_1$ to $t = t_2$, we perform a projective measurement $\{P_{(k)} = \sum_i |k, i_P\rangle \langle k, i_P|\}$ on the probe, where $\{|k, i_P\rangle\}$ are pure states of the probe. At $t = t_2$, we obtain a result k with probability p_k and the state of SB becomes

$$\rho_2 = \sum_k p_k \rho_2^{(k)} = \sum_{i,k} q_{i,k} \rho_2^{ik}, \quad (10)$$

where

$$\begin{aligned} \sqrt{q_{i,k}}\, |\psi^{ik}_{SBR}\rangle &= (\langle k, i_P| \otimes 1_{SBR}) |\psi_{PSBR}\rangle \quad (11) \\ \rho_2^{ik} &= {\rm tr}_R[|\psi^{ik}_{SBR}\rangle \langle \psi^{ik}_{SBR}|], \quad (12) \\ p_k \rho_2^{(k)} &= {\rm tr}_{PR}[\sum_i q_{i,k} |\psi^{ik}_{SBR}\rangle \langle \psi^{ik}_{SBR}|], \quad (13) \end{aligned}$$

with $|\psi^{ik}_{SBR}\rangle$ and $\rho_2^{(k)}$ being normalized. We can interpret the above as performing a measurement $\{M_{(k)}\}$, where

$$M_{(k)} = \sum_i \langle k, i_P| U_{SP} |0_P\rangle, \quad (14)$$

on S from $t = t_1$ to $t = t_2$. Then, the QC-mutual information [6, 8] is expressed as

$$I_{\rm QC} = S(\rho_1) - \sum_k p_k S(\rho_2^{(k)}). \quad (15)$$

We emphasize the following two points. First, we can determine the unitary interaction U_{SP} and the projective measurements $P_{(k)}$ for any measurement $M_{(k)}$. Hence, if we can evaluate the QC-mutual information $I_{\rm QC}$, then we can also evaluate $I_{\rm E}$. Second, the timings at which $I_{\rm E}$ and $I_{\rm QC}$ are defined are different. The information $I_{\rm E}$ is defined when only U_{SP} is completed, whereas the information $I_{\rm QC}$ is defined when the measurement $\{M_{(k)}\}$ is also completed. Thus, for two measurements with the same U_{SP}, $I_{\rm E}$ takes the same value but $I_{\rm QC}$ may take different values.

From $t = t_2$ to $t = t_3$, we perform a feedback control depending on the measurement result k. To be precise, we perform a unitary transformations $U_{(k)}$ on SB. At $t = t_3$, the state of SB is given by

$$\rho_3 \equiv \sum_k p_k U_{(k)} \rho_2^{(k)} U^\dagger_{(k)}. \quad (16)$$

From $t = t_3$ to $t = t_{\rm f}$, we choose a thermodynamic operation $U_{\rm fin}$ whose final state is equilibrium and perform it. Note that we only assume that the final state is macroscopically equilibrium; the final state may not be a canonical distribution given by

$$\rho_{\rm f}^{\rm can} \equiv \frac{\exp[-\beta' \hat{H}_{\rm f}^S]}{Z_{\rm f}^S(\beta')} \otimes \frac{\exp[-\beta_1 \hat{H}^{B_1}]}{Z^{B_1}(\beta_1)} \otimes ... \otimes \frac{\exp[-\beta_n \hat{H}^{B_n}]}{Z^{B_n}(\beta_n)}, \tag{17}$$

where β' is the inverse temperature of the final state of the system.

For the above process, we present the following four results and one conjecture:

Result 1: For any measurement $\{M_{(k)}\}$, the following inequality holds:

$$I_{\rm QC} \leq I_{\rm E}. \tag{18}$$

The inequality (18) allows us to substitute $I_{\rm E}$ for $I_{\rm QC}$ in any inequality of the form

$$x \leq y + zI_{\rm QC} \;\text{ or }\; u - vI_{\rm QC} \leq w, \tag{19}$$

where x, y, $z \geq 0$, u, $v \geq 0$ and w are real values. Thus, from the inequality [6]

$$\frac{U^S - F^S}{T} + \sum_{m=1}^{n} \frac{Q_m}{T_m} \leq -\frac{U'^S - F'^S}{T'} + k_B I_{\rm QC} \tag{20}$$

we obtain

$$\frac{U^S - F^S}{T} + \sum_{m=1}^{n} \frac{Q_m}{T_m} \leq -\frac{U'^S - F'^S}{T'} + k_B I_{\rm E}. \tag{21}$$

When the the system undergoes an isothermal process in contact with a single heat bath B at temperature T, the inequality (21) reduces to (2). Note that the substitution of $I_{\rm E}$ for $I_{\rm QC}$ may be valid for results of information thermodynamics other than (20).

Result 2: We can always achieve the new upper bound in the approximation such that the upper bound of the stadard second law $W_{\rm ext} \leq -\Delta F$ is always achievable.

Result 3: When the following conditions are satisfied, we can always achieve the equality of (21) with proper choices of $\{P_{(k)}\}$ and $\{U_{(k)}\}$ for any U_{SP}:

Condition 1: The system S is a two-level system.

Condition 2: The thermodynamic operations $U_{\rm init}$ and $U_{\rm fin}$ are strictly quasi-static.

Condition 3: $\vec{\lambda}^{SB}(t) = \vec{\lambda}_0$ is satisfied for $t_1 \leq t \leq t_2$.

Condition 2 corresponds to the condition for achieving the equality of (4), to which (21) reduces in a thermodynamic process without feedback control. Condition 3 implies that the system and baths do not interact during the measurement; if the system and baths interact during the measurement, the information obtained by the probe contains the information about the system as well as the baths. Thus, we can interpret Result 3 as follows: We can completely use the information obtained by the probe with a proper interpretation $\{P_{(k)}\}$ and a proper feedback $\{U_{(k)}\}$, if we do not waste energy during the thermodynamic processes and if the information describes only the system.

We conjecture that we can remove Condition 1:

Conjecture 1: Result 3 is valid without Condition 1.

Result 4: Under Conditions 1–3, there is a measurement $\{M_{(k)}\}$ for which we cannot achieve the equality of (20) with any $U_{(k)}$.

Results 1, 3 and 4 seem to contradict each other. Though the upper bound of (21) is always achievable and though inequality (18) exists, there is a case in which the upper bound of (20) is not achievable. However, the contradiction is spurious, not real. Note that when $I_{\rm E}$ is determined, we can take $\{P_{(k)}\}$ freely; in other words, we can choose the "best" interpretation of the information obtained by the probe. On the other hand, when $I_{\rm QC}$ is determined, $\{P_{(k)}\}$ is also determined already, and thus our interpretation of the probe's information is fixed uniquely.

Let us prove Results 1–3.

Proof of Result 1: Because of the definitions (8) and (15), we prove

$$\sum_k p_k S(\rho_2^{(k)}) \geq E_F^{SB\text{-}R}(\rho_{SBR}). \tag{22}$$

We can express the above as follows:

$$\begin{aligned} \sum_k p_k S(\rho_2^{(k)}) &= \sum_k p_k S(\sum_i \frac{q_{i,k}}{p_k} \rho_2^{(ik)}) \\ &\geq \sum_{i,k} q_{i,k} S(\rho_2^{(ik)}) = \sum_{i,k} q_{i,k} E^{SB\text{-}R}(|\psi_{SBR}^{ik}\rangle) \\ &\geq E_F^{SB\text{-}R}(\rho_{SBR}), \end{aligned} \tag{23}$$

where $\rho_{SBR} = \sum_{i,k} q_{i,k} \left|\psi_{SBR}^{ik}\right\rangle \left\langle\psi_{SBR}^{ik}\right| = {\rm tr}_P[|\psi_{PSBR}\rangle \langle\psi_{PSBR}|]$. □

Proof of Result 2: In the approximation, we can always achieve the upper bound of (20) for any measurement $\{M_{(k)}\}$ [7]. For an arbitrary unitary U_{SP}, there is a projective measurement $\{P_{(k)}\}$ that satisfies $I_{\rm E} = I_{\rm QC}$. Therefore, in the approximation, we can always achieve the upper bound of (21). □

Proof of Result 3: As in the derivation of (20) in Ref. [6], we can obtain the inequality (21) by transforming

$$S(\rho_{\rm i}) \leq -{\rm tr}[\rho_{\rm f} \log \rho_{\rm f}^{\rm can}] + I_{\rm E}. \tag{24}$$

Thus, we only have to prove that for any U_{SP}, we can always take $\{P_{(k)}\}$ and $\{U_{(k)}\}$ that satisfy

$$S(\rho_{\rm i}) = -{\rm tr}[\rho_{\rm f} \log \rho_{\rm f}^{\rm can}] + I_{\rm E}. \tag{25}$$

First, we prove that if ρ_3 in (16) is a canonical distribution, we can transform (25) into

$$E_F^{S\text{-}R}(\rho_{SR}) = S(\rho_3^S), \tag{26}$$

where $\rho_{SR} \equiv \mathrm{tr}_B[\rho_{SBR}]$, $\rho_3^S \equiv \mathrm{tr}_B[\rho_3]$, and $E_F^{S\text{-}R}$ is the entanglement of formation between S and R. Thanks to (8) and $S(\rho_\mathrm{i}) = S(\rho_1)$, we can transform (25) into

$$0 = -\mathrm{tr}[\rho_\mathrm{f} \log \rho_\mathrm{f}^\mathrm{can}] - E_F^{SB\text{-}R}(\rho_{SBR}). \tag{27}$$

Because of Condition 2, if ρ_3 is a canonical distribution, ρ_f is the canonical distribution $\rho_\mathrm{f}^\mathrm{can}$ in (17). Then we can transform (27) into

$$E_F^{SB\text{-}R}(\rho_{SBR}) = S(\rho_3), \tag{28}$$

where we use $S(\rho_3) = S(\rho_\mathrm{f})$. Thus, we only have to transform (28) into (26). If the following three equations hold, (28) and (26) are equivalent;

$$\begin{aligned} E_F^{SB\text{-}R}(\rho_{SBR}) &= E_F^{S\text{-}R_1}(\rho_{SR_1}) + E^{B\text{-}R_2}(|\psi_{BR_2}\rangle), && (29)\\ E_F^{S\text{-}R}(\rho_{SR}) &= E_F^{S\text{-}R_1}(\rho_{SR_1}), && (30)\\ S(\rho_3) &= S(\rho_3^S) + E^{B\text{-}R_2}(|\psi_{BR_2}\rangle), && (31) \end{aligned}$$

where we divide R into a two-level subsystem R_1 and the rest R_2.

Let us first prove (29). Owing to Condition 2 and the fact that ρ_i is a canonical distribution, ρ_1 is a canonical distribution as well. Because of Condition 3, $\hat{H}(t_1) = \hat{H}^S \otimes \hat{H}^B$ is valid. Thus, because of Condition 1, under the proper basis of R we can divide R into a two-level subsystem R_1 and a subsystem R_2 and express $|\psi_{SBR}\rangle$ as follows:

$$|\psi_{SBR}\rangle = |\psi_{SR_1}\rangle \otimes |\psi_{BR_2}\rangle\,. \tag{32}$$

Owing to (7) and (32), we can express $|\psi_{PSBR}\rangle$ as follows: $|\psi_{PSBR}\rangle = |\psi_{PSR_1}\rangle \otimes |\psi_{BR_2}\rangle$. Thus, we can express ρ_{SBR} as

$$\rho_{SBR} = \rho_{SR_1} \otimes |\psi_{BR_2}\rangle\langle\psi_{BR_2}|\,, \tag{33}$$

where $\rho_{SR_1} \equiv \mathrm{tr}_P[|\psi_{PSR_1}\rangle\langle\psi_{PSR_1}|]$. From (33), we have (29) and (30). Next, we prove (31). Note that S has been isolated for $t_1 \le t \le t_3$ with proper $\{U_{(k)}\}$. We can therefore express ρ_3 as $\rho_3^S \otimes |\psi'_{BR_2}\rangle\langle\psi'_{BR_2}|$ with $E^{B\text{-}R_2}(|\psi_{BR_2}\rangle) = E^{B\text{-}R_2}(|\psi'_{BR_2}\rangle)$. Thus, we have (31).

Now, we only have to find $\{P_{(k)}\}$ on P and $\{U_{(k)}\}$ on S such that ρ_3 and ρ_3^S are canonical distributions and that (26) holds. We first prove that if ρ_3^S is a canonical distribution, ρ_3 is also a canonical distribution. To prove this, we only have to note that $\mathrm{tr}_{R_2}[|\psi'_{BR_2}\rangle\langle\psi'_{BR_2}|]$ is a canonical distribution because B has been isolated for $t_1 \le t \le t_3$ and because $\mathrm{tr}_S[\rho_1]$ is a canonical distribution. We second find $\{P_{(k)}\}$ on P and $\{U_{(k)}\}$ on S such that ρ_3^S is a canonical distribution and that (26) holds. Because both S and R_1 are two-level systems, we can treat the state $|\psi_{PSR_1}\rangle$ as a three-qubit pure state under a proper basis of P. In the supplementary materials, we prove the following with the approach used in Ref. [22]: we can perform a projective measurement $\{\tilde{P}_{(k)}\}_{k=0,1}$ on the probe P such that the results $\tilde{P}_{(k)}|\psi_{PSR_1}\rangle$ are LU-equivalent for $k = 0, 1$ and $E^{S\text{-}R_1}(\tilde{P}_{(k)}|\psi_{PSR_1}\rangle) = E_F^{S\text{-}R_1}(\rho_{SR_1})$ is valid. Because the results $\tilde{P}_{(k)}|\psi_{PSR_1}\rangle$ are LU-equivalent, there exists $\{V_{(k)}\}_{k=0,1}$ on S, which satisfies $\rho_3^\mathrm{pre} \equiv V_{(k)}\rho_2^{S(k)}V_{(k)}^\dagger$, where $\rho_2^{S(k)} \equiv \mathrm{tr}_{R_1}[\tilde{P}_{(k)}|\psi_{PSR_1}\rangle\langle\psi_{PSR_1}|\tilde{P}_{(k)}]$. Owing to Condition 1, if $E^{S\text{-}R_1}(\tilde{P}_{(k)}|\psi_{PSR_1}\rangle) \neq 0$, we can make the state $\rho_3^S = \hat{V}\rho_3^\mathrm{pre}\hat{V}^\dagger$ a canonical distribution with a unitary transformation V on S. Thus, $\{\tilde{P}_{(k)}\}$ and $\{VV_{(k)}\}$ are the measurement and feedback that we want.□

Proof of Result 4: It is sufficient to prove the existence of a counterexample of the measurement $\{M_{(k)}\}$. The equality of (20) is valid only if there exists a set of unitary transformations $\{U_{(k)}\}$ that satisfy $\sum_k p_k S(\rho_2^{(k)}) = S(\rho_3)$ [6]. We can transform $S(\rho_3)$ as follows:

$$\begin{aligned} S(\rho_3) &= S(\sum_{k=0,1} p_k U_{(k)}\rho_2^{(k)}U_{(k)}^\dagger) \\ &= \sum_{k=0,1} p_k S(\rho_2^{(k)}) + \sum_{k=0,1} p_k D(U_{(k)}\rho_2^{(k)}U_{(k)}^\dagger||\rho_3), \end{aligned} \tag{34}$$

where $D(\rho||\rho') \equiv \mathrm{tr}[\rho(\log\rho - \log\rho')]$. Because $D(\rho||\rho') = 0$ if and only if $\rho = \rho'$ and because of (34), the equation $\sum_k p_k S(\rho_2^{(k)}) = S(\rho_3)$ is valid if and only if $\{\rho_2^{(k)}\}$ are LU-equivalent for $k = 0, 1$, in other words, if and only if the measurement $\{M_{(k)}\}$ is a deterministic measurement. Based on this logic, if Result 4 were not valid, any measurement $\{M_{(k)}\}$ would be deterministic. This is clearly false, and thus Result 4 holds.□

To conclude, we obtain a new version of information thermodynamic inequality. In this inequality, the information gain is the entanglement gain; the new information I_E represents the amount of the entanglement between the system and the response system which the probe takes from the system. The new information depends only on the premeasurement state of the system and the unitary interaction between the probe and the system, and thus when I_E is determined, we can take $\{P_{(k)}\}$ freely. The QC-mutual information I_QC does not have this freedom; Results 3 and 4 follow from this difference of the freedom between I_E and I_QC. Thus, in the above context, we can say as follows: the substance of information is the entanglement. The information gain is already completeded when the unitary interaction is over and the projective measurement is only the interpretation of the information.

This work was supported by the Grants-in-Aid for Japan Society for Promotion of Sci- ence (JSPS) Fellows (Grant No. 24E8116). The author thanks Prof. Naomichi Hatano for useful discussions.

SUPPLEMENTARY INFORMATION

In the present supplementary material, we prove the following theorem:

Theorem 1 *For an arbitrary three-qubit pure state* $|\psi_{PSR_1}\rangle$*, there exists a projective measurement* $\{P_{(k)}\}_{k=0,1}$ *such that* *the results* $P_{(k)}\left|\psi_{PSR_1}\right\rangle$ *are LU-equivalent for* $k=0,1$ *and*

$$E^{S\text{-}R_1}(P_{(k)}\left|\psi_{PSR_1}\right\rangle)=E_F^{S\text{-}R_1}(\rho_{SR_1}) \tag{35}$$

is valid.

Proof: Because ρ_{SR_1} is a two-qubit mixed state, we can express $E_F^{S\text{-}R_1}(\rho_{SR_1})$ in the form of the concurrence [23]:

$$E_F^{S\text{-}R_1}(\rho_{SR_1})=h\left(\frac{1+\sqrt{1-C^2_{SR_1}(\rho_{SR_1})}}{2}\right), \tag{36}$$

where $C_{SR_1}(\rho_{SR_1})$ is the concurrence of ρ_{SR_1} and $h(x)\equiv -x\log x-(1-x)\log(1-x)$. Thus, we only have to find a projective measurement $\{\tilde{P}_{(k)}\}_{k=0,1}$ such that $\tilde{P}_{(k)}\left|\psi_{PSR_1}\right\rangle$ are LU-equivalent for $k=0,1$ and $C_{SR_1}(\tilde{P}_{(k)}\left|\psi_{PSR_1}\right\rangle)=C_{SR_1}(\rho_{SR_1})$.

Before giving the projective measurement $\{\tilde{P}_{(k)}\}$, we first present preparations. First we express $\left|\psi_{PSR_1}\right\rangle$ in the form of the generalized Schmidt decomposition [24]:

$$\begin{aligned}\left|\psi_{PSR_1}\right\rangle &= \lambda_0\left|000\right\rangle+\lambda_1 e^{i\varphi}\left|100\right\rangle+\lambda_2\left|101\right\rangle \\ &\quad +\lambda_3\left|110\right\rangle+\lambda_4\left|111\right\rangle \quad (0\leq\varphi\leq\pi)\end{aligned} \tag{37}$$

and introduce the following eight parameters [22];

$$\begin{aligned}
K_{PS} &\equiv C^2_{PS}+\tau_{PSR_1}, &(38)\\
K_{PR_1} &\equiv C^2_{PR_1}+\tau_{PSR_1}, &(39)\\
K_{SR_1} &\equiv C^2_{SR_1}+\tau_{PSR_1}, &(40)\\
J_5 &\equiv 4\lambda_0^2(|\lambda_1\lambda_4 e^{i\varphi}-\lambda_2\lambda_3|^2+\lambda_2^2\lambda_3^2-\lambda_1^2\lambda_4^2), &(41)\\
K_5 &\equiv J_5+\tau_{PSR_1}, &(42)\\
\Delta_J &\equiv K_5^2-K_{PS}K_{PR_1}K_{SR_1}, &(43)\\
e^{-i\tilde{\varphi}_5} &\equiv \frac{\lambda_2\lambda_3-\lambda_1\lambda_4 e^{i\varphi}}{|\lambda_2\lambda_3-\lambda_1\lambda_4 e^{i\varphi}|}, &(44)\\
Q_{\mathrm{e}} &\equiv \mathrm{sgn}\left[\sin\varphi\left(\lambda_0^2-\frac{\tau_{PSR_1}+J_5}{2(C^2_{SR_1}+\tau_{PSR_1})}\right)\right], &(45)
\end{aligned}$$

where τ_{PSR_1} is the tangle of $\left|\psi_{PSR_1}\right\rangle$ and sgn$[x]$ is the sign function,

$$\mathrm{sgn}[x]=\left\{\begin{array}{ll} x/|x| & (x\neq 0)\\ 0 & (x=0)\end{array}\right\}. \tag{46}$$

When $Q_{\mathrm{e}}=0$, there are two possible decompositions which satisfy (37). We then choose the decomposition with a greater coefficient λ_0.

Now the preparation has been completed. In the basis of (37), the projective measurement $\{\tilde{P}_{(k)}\}_{k=0,1}$ is given as follows;

$$\tilde{P}_{(0)}=\sqrt{\begin{pmatrix} a & ke^{-i\theta}\\ ke^{i\theta} & b\end{pmatrix}},\ \tilde{P}_{(1)}=\sqrt{\begin{pmatrix} 1-a & -ke^{-i\theta}\\ -ke^{i\theta} & 1-b\end{pmatrix}} \tag{47}$$

where the measurement parameters a, b, k and θ are defined as follows:

$$\begin{aligned}
a &= \frac{1}{2}-\frac{K_5\tau_{PSR_1}\dot{\pm}\sqrt{\Delta_J}C^2_{SR_1}}{2K_{SR_1}\sqrt{K_5^2-K_{PS}K_{PR_1}C^2_{SR_1}}}, &(48)\\
b &= 1-a, &(49)\\
k &= \sqrt{a(1-a)}, &(50)\\
\theta &= -\tilde{\varphi}_5, &(51)
\end{aligned}$$

and when $Q_{\mathrm{e}}\neq 0$ the mark $\dot{\pm}$ means $-Q_{\mathrm{e}}$ and when $Q_{\mathrm{e}}=0$ the mark $\dot{\pm}$ means $-$. With using (48)–(51) and Lemma 1 of Ref. [22] and after straightforward algebra, we can confirm that the measurement $\{\tilde{P}_{(k)}\}$ is the measurement that we sought.

[1] L. Szilard, Z. Phys. **53**, 840 (1929).
[2] R. Landauer, IBM J. Res. Dev. **5**, 183 (1961).
[3] C. H. Bennett, Int. J. Theor. Phys. **21**, 905 (1982).
[4] B. Piechocinska, Phys. Rev. A **61**, 062314 (2000).
[5] M. A. Nielsen, C. M. Caves, B. Schumacher, and H. Barnum, Proc. R. Soc. A **454**, 277 (1998).
[6] T. Sagawa and M. Ueda, Phys. Rev. Lett, **100** 080403 (2008).
[7] K. Jacobs, Phys. Rev. A **80**, 012322 (2009).
[8] T. Sagawa and M. Ueda, Phys. Rev. Lett. **102**, 250602 (2009).
[9] F. J. Cao and M. Feito, Phys. Rev. E **79**, 041118 (2009).
[10] S. Toyabe, *et al.*, Nature Phys. **6**, 988 (2010).
[11] J. M. Horowitz and J. M. R. Parrondo, New J. Phys. **13**, 123019 (2011).
[12] S. D. Liberato and M. Ueda, Phys. Rev. E **84**, 051122 (2011).
[13] D. Deutsch, Proc. R. Soc. Lond. A **400**, 97 (1985)
[14] C. H. Bennett, *et al.*, Phys. Rev. Lett. **70**, 1895 (1993)
[15] C. H. Bennett and S. J. Wiesner, Phys. Rev. Lett. **69**, 2881 (1992)
[16] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information (Cambridge University Press, Cambridge, 2000)
[17] C. H. Bennett, *et al.*, Phys. Rev. A **53**, 2046 (1996).
[18] C. H. Bennett, *et al.*, Phys. Rev. A **54**, 3824 (1996).
[19] A. Kitaev and J. Preskill, Phys. Rev. Lett. **96**, 110404 (2006).
[20] G. Vidal, Phys. Rev. Lett. **99**, 220405 (2007).
[21] Sandu Popesucu, *et al.*, Nature Phys. **2**, 754 (2006)
[22] H. Tajima, Ann. of Phys. (2012), doi:10.1016/j.aop.2012.11.001
[23] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
[24] A. Acín, *et al.*, Phys. Rev. Lett. **85** 1560 (2000).